

# Differences between SNAGS and Mc Kenzie Interventions with Core Stability and Mc Kenzie on the Functional Ability of Elderly People with Lumbar HNP Conditions at Hermina Hospital Depok

Irma Megawanty<sup>1\*</sup>, Luluk Maulina<sup>2</sup>

<sup>1,2</sup> Program Studi Sarjana Fisioterapi, Fakultas Ilmu Kesehatan, Universitas Medika Suherman, Indonesia

Email: [irmamegawanty11@gmail.com](mailto:irmamegawanty11@gmail.com)

Submission : 2023-07-29; Accepted : 2023-11-06; Published : 2023-11-19

## ABSTRACT

**Introduction:** HNP is a degenerative disease where the elasticity of the annulus fibrosus decreases so that it becomes one of the causes of LBP and often occurs in the elderly. One of the roles of physiotherapy in HNP cases can be treated with exercise therapy, with the SNAGS and Mc Kenzie intervention methods as well as with Core Stability and Mc Kenzie. To help improve the functional abilities of the elderly. The purpose of the study is to determine the differences in interventions for Lumbar HNP cases using the SNAGS and Mc Kenzie methods as well as Core Stability and Mc Kenzie on the functional abilities of the elderly. **Methods:** The design of this study used a quasi-experimental with two groups pre-post test. A sample of 22 people was selected by purposive sampling which was divided into two, namely treatment 1 and treatment 2. The research was conducted for 4 weeks consisting of one pretest, with meetings twice a week with a duration of 30-45 minutes and one post-test. The instrument used is the ODI questionnaire. **Results:** The result showed that the normality test using the Shapiro-Wilk test resulted in a value of  $p=0.000$ , which means it was significant in both treatment groups. From the results of the paired test, it is significant with a value of  $p = 0.000$  meaning  $p < 0.05$ . The reduction of the difference in the percentage of disability in treatment group 1 was -32.55 per cent, and in treatment group 2 was -37.27 per cent. **Conclusion:** The conclusion from the study is SNAGS and Mc Kenzie interventions as well as Core Stability and Mc Kenzie can affect the functional abilities of the elderly. With more effective results on Core stability and Mc Kenzie interventions in improving the functional abilities of the elderly with Lumbar HNP conditions.

**Keywords:** HNP, SNAGS, Mc Kenzie, Core Stability, Elderly

ISSN 2722 – 9610  
 E –ISSN 2722 - 9629

## INTRODUCTION

Health Law No. 36 of 2009 is a government effort to explain the meaning of health, namely mentally, physically and socially so that you can do things that are productive and economical. Health is the right of all people, including the elderly (Sipayung, Anggiat and Soeparman, 2020)

UU no. 13 of 1998 explains that it is considered elderly if a person is over 60 years old (Ministry of Health of the Republic of Indonesia, 2013). At this age, humans will experience various changes in all aspects of life, especially health, including biological, sexual, emotional,

social and spiritual changes. (Wijoyo & Daulima, 2020).

One of the studies regarding the welfare of the elderly (men and women) aged 60 years and over who experienced a natural decline in physical and psychological abilities which resulted in slowed movements, shorter footsteps, and decreased muscle strength, especially in the lower limbs. Reduced leg strength means that you cannot tread firmly and are unstable, so steps must be taken to anticipate balance problems such as slipping and tripping. (Sipayung, Anggiat and Soeparman, 2020)



Jumping, running, exercising and walking normally are difficult things for the elderly because their health, especially their physical health, is declining. (Ariyanto et al., 2020).

In the elderly, namely, those aged 60-70 years, pathological processes are often found that cause a decrease in balance and deviations in gait, one of which is characterized by a slowed walking pattern. Changing walking patterns in the elderly is an effort to improve balance. Limited activity and abnormal motor skills are caused by impaired balance whose function decreases so elderly people often use assistive devices or help from other people in carrying out their activities. Improving health, and physical and social quality will certainly increase life expectancy. According to WHO, there will be an increase of  $\pm$  600,000,000 to 2,000,000,000 elderly people by 2050. The Asian region will have an elderly population that will increase by 82 per cent in the next 25 years (Wijayani et al., 2022).

A total of 9.03 per cent or 23.66 million people is the projected data for elderly people in Indonesia in 2017. The elderly population in 2020 is estimated to reach 27.08 million, in 2025 (33.69 million), and in 2030 (40, 95 million), while in 2035 it could reach 48.19 million. The elderly have a decline in physical and psychological development which leads to negative, causing dependency (Adi et al., 2020).

Problems that often occur in the elderly include not having access to education, health, old age security, social support from family and friends and physical exercise (Ariyanto et al., 2020).

Disorders that occur in the elderly include 25 per cent needing hospitalization for further evaluation, 20 per cent of sufferers of which 13 per cent need medical assistance and 40 per cent experience sleep disorders (Nugroho et al., 2018).

Low back pain affects 80 per cent of the population once in a lifetime. According to health workers in Indonesia, the prevalence of musculoskeletal disease in Indonesia is 11.9 per cent based on diagnosis and 24.7 per cent with symptomatic diagnosis, while the highest prevalence of the musculoskeletal disease is 31.2 per cent for those who work as farmers, fishermen or labourers. At the age of 35-55 years, there will be a decline and abnormalities in the

intervertebral discs so the risk of lower back pain will be higher (Rikesdas, 2013).

One of the causes of lower back pain is Herniated Nucleus Pulposus (HNP), which means a long-term condition of protrusion of the intervertebral disc caused by injury and incorrect mechanical load. However, the main cause of HNP is the tearing of the annulus fibrosus which results in decreased elasticity due to degeneration (Widyasari & Wulandari, 2020).

Complaints that are felt if HNP occurs are decreased muscle strength which interferes with the patient's activities, limited joint movement, pain and tingling in the lower waist to the legs, and spasms (Dwi et al., 2020).

The causes of HNP include degenerative changes which cause decreased flexibility and thinning of the nucleus pulposus, and moderate but repeated trauma involving the intervertebral disc which results in tearing of the annulus fibrosus (Widyasari & Wulandari, 2020).

Based on estimates, 40 per cent of the Indonesian population at the age of 65 years have experienced complaints in the waist with a prevalence in men of 18.2 per cent and women of 13.6 per cent, while the incidence of HNP from patient visits to hospitals is in the range of 3-17 per cent. After headaches, the complaint that ranks second is HNP (Sani & Durahim, 2021).

Based on the survey results at Hermina Hospital, there were 24 respondents as samples, but 2 respondents did not meet the research requirements so the total research sample was 22 elderly patients experiencing Lumbar HNP from December 2022-January 2023. Incidents of lumbar disability experienced by disabled patients when carrying out activities involve lowering movements such as sitting, standing and walking (RS Hermina Depok, 2022).

With the problems experienced by the elderly, the development of physiotherapy treatment emerged (Dwi et al., 2020).

Physiotherapy is a form of health service intended for individuals or groups to develop, maintain and restore body movement and function throughout the life cycle using manual treatment, electrical equipment (electrotherapy and mechanical), functional training and communication. (Indonesian Ministry of Health, 2013). Another opinion states that physiotherapy



is a provider of health services for individuals and groups to develop, maintain and restore maximum movement and functional abilities throughout the life cycle (Sipayung et al., 2020).

Physiotherapy is an active form of treatment that focuses on joint and muscle movement. (Sipayung et al., 2020).

In sufferers of LBP due to HNP, exercise therapy is carried out as a form of physiotherapy (Sipayung, Anggiat and Soeparman, 2020).

The benefits of exercise therapy in the form of body movements, postures and other activities include: correcting or preventing disorders, improving, restoring or increasing physical function, preventing or reducing health-related risk factors, and optimizing health conditions, fitness or overall sense of well-being. (Amin et al., 2018).

To reduce movement and function disorders in the elderly, you can use the Mc Kenzie method to restore the existing nucleus pulposus and reinsert it into the annulus pulposus (Sipayung, Anggiat and Soeparman, 2020).

Mc Kenzie is a useful exercise for strengthening the back muscles and relaxing the abdominal muscles, strengthening the transverse abdominis, internal obliquus and external obliquus muscles. (Maycahyasari et al., 2022). Mc Kenzie has the characteristic that posture correction ensures body posture is in the right condition (Maycahyasari et al., 2022).

Brian R. Mulligan, a physiotherapist from New Zealand, introduced manual therapy for Spinal sustained natural apophyseal glides (SNAGS) on the lumbar vertebrae which has the characteristic of mobilization with movement (MWM) or a combination of accessory passive movements during active or physiological passive movement techniques. SNAGS is nonoscillatory mobilization which has been implemented on the lumbar vertebrae which is an MWM procedure. SNAGS has been widely applied in orthopaedic manual physical therapy (OMPT) because in vertebral conditions this technique has been declared effective (Sani & Durahim, 2021).

To produce optimal movement when carrying out activities, it is necessary to increase the trunk muscle groups around the rear driver with Core Stability exercises (Tantangan et al., 2021). This aims to improve neuromuscular

control, endurance, and muscle strength so that the spine is stable and dynamic as shown in research with chronic low back pain. (Harder, 2021). Good core stability can function to increase movement and prevent injury because it is an important factor in body posture. In medical literature, core stability is 'the product of motor control and muscle capacity in the lumbar, pelvic, hip complex', in musculoskeletal terms including the spine, pelvis and hip joints, as well as the proximal lower extremities in addition to all related muscles. (Zulvikar, 2016).

The general aim of this research is to find out whether there is a difference in administering SNAGS and Mc Kenzie measures to the functional abilities of elderly people with Lumbar HNP conditions and to find out whether there are differences in giving Core Stability & Mc Kenzie measures to the functional abilities of elderly people with Lumbar HNP conditions. Meanwhile, the specific objective is to determine the actions of SNAGS & Mc Kenzie on increasing the functional abilities of elderly people with Lumbar HNP conditions and to find out the actions of Core Stability & Mc Kenzie on increasing the functional abilities of elderly people with Lumbar HNP conditions. It is hoped that this research will become a means of increasing the highest level of health for all levels of society as a whole. Apart from that, as a reference for knowing SNAGS actions, Mc. Kenzie and Core Stability on the functional abilities of elderly people with Lumbar HNP conditions and increasing knowledge of exercise techniques as actions based on physiotherapy science. So that in further research we can find out and understand about the elderly and the condition of Lumbar HNP and prove whether SNAGS, Mc Kenzie & Core Stability are effective for the functional abilities of elderly people with Lumbar HNP conditions.

## METHODS

*Quasi-experimental with a two-group pretest and posttest design* is the type of research used in this study using 2 objects, namely treatment group 1 which was given SNAGS & Mc Kenzie actions, while treatment group 2 was given Core Stability & Mc Kenzie actions. The instruments required in the research consisted of an *informed concern* form, research explanation



form, ODI questionnaire, respondents, stationery and a bed for carrying out exercise therapy.

The sample used 24 respondents as a sample, but 2 respondents did not meet the research requirements so the total research sample was 22 elderly patients who were divided into 2 groups, namely 11 people in treatment group 1, namely snags and MC Kenzie training and 11 people in treatment group 2, namely core training stability and MC Kenzie at RS Hermina Depok. The sampling technique was simple random sampling with the inclusion criteria of elderly men and women aged over 60 years, willing to take part in the entire research program, able to carry out SNAGS, Core Stability & Mc. Kenzie's actions, in Lumbar HNP conditions. Data collection through pre-test, research implementation and post-test. Sample exclusion criteria dropped out due to stopping therapy. Data processing includes editing, coding, processing and cleaning. Data analysis technique using graphpad prism. First, a normality test was carried out using the Shapiro-Wilk test, then a hypothesis test was carried out, namely the Paired t-test with a significance level of  $p < 0.05$ . The final stage is interpretation and drawing conclusions based on the research results and analysis obtained.

## RESULTS

Table 1. Distribution of Research Samples Based on Age and Gender

Variabel	Kategori	Kelompok perlakuan 1		Kelompok perlakuan 2	
		N	%	n	%
Usia (tahun)	60-70	5	45%	3	27%
	70-80	6	55%	8	73%
Total		11	100%	11	100%
Mean ± SEM		68,9±0,431		71,7±0,373	
Jenis	Laki-laki	3	27%	2	18%
Kelamin	Perempuan	8	73%	9	82%
Jumlah		11	100%	11	100%

Based on Table 1, the age characteristics of this research sample ranged from 60-80 years. In treatment group 1, the largest sample age was between 70-80 years, there were 6 samples (55 per cent), and in 60-70 years, there were 5 samples (45 per cent). In treatment group 2, the largest sample age was also between 70-80 years, there were 8 samples (73 per cent), 60-70 years old, and there were 3 samples (27 per cent). In

terms of gender, it was found that women dominated, as shown in treatment group 1, the sample size was 8 female respondents (73 per cent) and 3 male respondents (27 per cent). Meanwhile, in treatment group 2, there were 9 female respondents (82 per cent) and 2 male respondents (18 per cent).

Table 2. ODI (Oswestry Disability Index) pre-test and post-test scores for Treatment Group 1 and Treatment Group 2.

Sampel	Kelompok perlakuan 1		Kelompok perlakuan 2	
	Pre test	Post test	Pre test	Post test
Mean±SEM	61,64%±0,451	29,09%±1,127	64,45%±0,627	27,18%±1,298

Based on the data in Table 2, shows that the post-test score for treatment group 1 has a mean value of 29.09 per cent with an SEM of 1.127, while the pre-test score obtained a mean value of 61.64 per cent with an SEM of 0.451. This is related to an increase in functional ability after administering the action. within 4 weeks.

In treatment group 2, it shows that the post-test score in treatment group 2 has a mean value of 27.18 per cent with an SEM of 1.298, while the pre-test score obtained a mean value of 64.45 per cent with an SEM of 0.627. This is related to an increase in functional ability after giving the action for 4 weeks.

Table 3. Data normality test with Shapiro Wilk test, pre-test and post-test in treatment group 1 and treatment group 2.

Kelompok	Shapiro Wilk Test		Distribusi Data
	P value		
	Pre test	Post test	
Perlakuan 1	0,5584	0,2630	Normal
Perlakuan 2	0,3602	0,8054	Normal

Based on the results of the Shapiro-Wilk normality test, the process uses a computerized system using the GraphPad Prism program. Table 3 shows that for each treatment group, both pre-test and post-test, the p-value obtained in treatment group 1, pre-test = 0.5584 and post-test = 0.2630. In treatment group 2 pre test = 0.3602 and post test = 0.8054. From the results of this data,  $p > 0.05$ , which means the data distribution is normal, so it is continued with the paired t-test.



Table 4. Paired T Test results for hypothesis testing on functional abilities using treatment group 1 and treatment group 2

Paired t Test			
Kelompok	Mean	Sd	P Value
Perlakuan 1	-32,55	11,78	0,000
Perlakuan 2	-37,27	9,678	

The data is presented in the form of mean and standard deviation and then hypothesis testing is carried out using a paired t-test with a significance of  $p < 0.05$ .

Based on the results of data analysis in each treatment group 1 and 2 using a paired t-test to see the functional ability of elderly people with Lumbar HNP conditions. Based on the results of the paired t-test analysis, the results for treatment group 1 were  $p = 0.000$  ( $p < 0.05$ ), which means  $H_a$  was accepted,  $H_o$  was rejected, then treatment group 2 was  $p = 0.000$  ( $p < 0, 05$ ) which means  $H_a$  is accepted and  $H_o$  is rejected, so it can be concluded that there is a significant increase in the functional ability of elderly people with Lumbar HNP conditions.

Based on the results of the hypothesis test, it was concluded that the two treatment groups had an increasing effect on the functional abilities of the elderly in Lumbar HNP conditions after being given treatment for 4 weeks. The effectiveness of treatments 1 and 2 can be seen by the mean difference between the two groups. The average difference value (mean) is -32.55 for treatment group 1 and the average difference value (mean) is -37.27 for treatment group 2, so it can be concluded that increasing the functional ability of elderly people who experience Lumbar HNP is a more effective treatment intervention 2 (Core Stability & Mc Kenzie) than treatment action 1 (SNAGS & Mc Kenzie).

## DISCUSSION

In this study, changes were found in the ODI (Oswestry Disability Index) scores in treatment groups 1 and 2.

Treatment group 1 experienced a change in the ODI (Oswestry Disability Index) score with the pre-test score being an average of 61.64 per cent and the post-test score being an average of 29.09 per cent. This means there is an increase in the functional ability of the elderly in Lumbar HNP conditions. The results of this research are

by research from (Sani & Durahim, 2021). The results of the research are that the administration and actions of SNAGs & Mc Kenzie influence reducing disability in Lumbar HNP conditions. This is a result of increased functional ability so that joint and muscle movements experience increased strength.

Meanwhile, in treatment group 2 there was a change in the ODI (Oswestry Disability Index) score with the pre-test score being an average of 64.45 per cent and the post-test score being an average of 27.18 per cent. This means there is an increase in the functional ability of the elderly in Lumbar HNP conditions. These results are in line with research (Tantangan et al., 2021) which reveals that these two methods are equally effective in increasing functional ability and can reduce lumbar HNP because these two methods impact the stabilizer muscles in the spine to become strong and relaxed, posture The body's faults are repaired so that joint pressure and ligament stretching are reduced, which can increase analgesic function, which is the function of the  $\beta$ -endorphin hormone. By carrying out regularly using the Core Stability & Mc Kenzie method, HNP sufferers will relax and improve their back muscles

This exercise can have an impact on not disrupting social life and functional activities by improving lower back disability in Lumbar HNP sufferers. (Zahratur & Priatna, 2019).

Based on the results of the normality test using the Shapiro-Wilk test, from the analysis of each treatment group, both pre-test and post-test, the p-value for treatment group 1, pre-test = 0.5584 and post-test = 0.2630. In treatment group 2 pre test = 0.3602 and post test = 0.8054. From the results of this data,  $p > 0.05$ , which means the data is distributed normally, so it is continued with the paired t-test.

From testing hypothesis 1 on data from treatment group 1, the results showed an increase in the functional ability of elderly people with Lumbar HNP conditions, using the paired t-test with a value of  $p = 0.000$ , meaning  $p < 0.05$  (smaller than 0.05), which means  $H_a$  is accepted and  $H_o$  is rejected, which means there is an influence of SNAGS and Mc Kenzie actions on the functional abilities of elderly people with Lumbar HNP conditions.



From testing hypothesis 2 on data from treatment group 2, the results showed an increase in the functional ability of elderly people with Lumbar HNP conditions, using the paired t-test with a value of  $p=0.000$ , meaning  $p<0.05$  (smaller than 0.05), which means  $H_a$  is accepted and  $H_o$  is rejected, which means that there is an influence of Core Stability & Mc Kenzie actions on the functional abilities of elderly people with Lumbar HNP conditions.

The effectiveness that occurred in treatment groups 1 and 2 can be seen from the difference in mean values in the two groups. The average difference value (mean) is -32.55 for treatment group 1 with the average difference value (mean) of -37.27 for treatment group 2, which means the action of treatment group 2 (Core Stability & Mc. Kenzie) is more effective than the treatment group 1 (SNAGS & Mc. Kenzie) in increasing the functional ability of elderly people in Lumbar HNP conditions.

The results of this research are in line with the theory of (Norlinta et al., 2019) that the Mulligan SNAGS technique is a manual therapy action by determining the facet joint where the injury occurs by applying passive gliding in the lumbar region so that it can increase functional and muscle strength. This technique will appear placebo (comfortable) because it is a combination of co-contraction movements and movement control without pain.

Based on the theory of (Tantangan et al., 2021) which is in line with the results of this research, providing Core Stability training in Lumbar HNP actions, to activate m. transfers abdominis and m. lumbar multifidus. These muscles act as the main stabilizers in the lumbar spine which play a role in balancing between activated agonist and antagonist muscles. When balance is achieved in the contraction of the abdominal muscles and the movement of the lumbar muscles when carrying out activities, it has an impact on increasing body awareness and correcting good posture due to adequate control of lumbar movement.

According to the theory of (Sipayung et al., 2020), McKenzie exercises are extension movements that dominate back exercises. This exercise uses the principle of stabilizing muscles in the spine by strengthening and relaxing. To

improve the functional abilities of elderly people with LBP due to HNP, the Mc Kenzie method can be used.

## CONCLUSION

Based on the research results, conclusions can be drawn, including: 1. SNAGS and Mc Kenzie's actions improve the functional abilities of elderly people with Lumbar HNP conditions. 2. Core Stability and Mc Kenzie's actions improve the functional abilities of elderly people with Lumbar HNP conditions. 3. There is a difference in the decrease in ODI scores between the two treatment groups, with the effectiveness value of treatment group 2 being more effective than treatment group 1.

## ACKNOWLEDGEMENT

Thank you to God Almighty for the smoothness provided during this research as well as the lecturers who always provided guidance in the implementation and preparation of this research as well as friends and family who always provided support for this research.

## REFERENCES

- Adi, S., Kurniawan, S., & Ardian, K. (2020). Efek Senam Lanjut Usia Terhadap Peningkatan Aktifitas Fisik Lanjut Usia Di Dusun Bonorejo Kelurahan Plesungan. *Wiraraja Medika : Jurnal Kesehatan*, 10(1), 17–22.
- Amin, A. A., Abidin, Z., & Widianingrum, U. (2018). Pengaruh Infra-Red, Tens Dan Low Back Core Stabilization Exercise Pada Kondisi Myalgia. *Jurnal Fisioterapi Dan Rehabilitasi*, 2(1), 17–25. <https://doi.org/10.33660/jfrwhs.v2i1.43>
- Ariyanto, A., Puspitasari, N., & Utami, D. N. (2020). Aktivitas fisik terhadap kualitas hidup pada lansia Physical Activity To Quality Of Life In The Elderly. *Jurnal Kesehatan Al-Irsyad*, XIII(2), 145–151.
- Dwi, W., Mu'jizatillah, & Fauziah, E. (2020). Management of Physiotherapy for Lumbar Functional Disorders due to Hernia Nucleus Pulposus with PNF Technique, TENS and McKenzie Exercise at RSUD Ulin Banjarmasin 2019. *Jurnal Kajian Ilmiah Kesehatan Dan Teknologi*, 2(1), 6–14.
- Harder, L. (2021). The effects of the McKenzie



- Method and strengthening/stabilization exercises in patients with chronic nonspecific low back pain. *Honors College Theses*.
- Kemendes RI No. 80 Tahun 2013. (2013). *Keputusan Menteri Kesehatan No.80 Tahun 2013*. 1536, 3. file:///C:/Users/User/Downloads/Documents/bn1536-2013.pdf
- Maycahyasari, D., Herawati, I., Hidayati, A., & Rahman, F. (2022). Penatalaksanaan Fisioterapi Pada Kasus Lansia Dengan Penurunan Fungsional Low Back Pain Di Klinik Atik Hidayati Davini. *Jurnal Kesehatan Dan Masyarakat*, 2(April), 13–18.
- Norlinta, S. N. O., Kuswardhani, R. T., Imron, M. A., Tirtayasa, K., Adiputra, N., & Irfan, M. (2019). Teknik Mulligan Paling Efektif Dibandingkan Mobilisasi Segmental Untuk Peningkatan Kemampuan Fungsional Pada Pembatik Tulis Dengan Cervical Syndrome. *Sport and Fitness Journal*. <https://doi.org/10.24843/spj.2019.v07.i02.p03>
- Nugroho, F., Weta, I. W., -, S., Griadhi, I. P. A., Satriyasa, B. K., & Irfan, M. (2018). Penambahan Nerve Stretching Lebih Baik Dibandingkan Nerve Gliding Setelah Mc Kenzie Exercise Dalam Menurunkan Gangguan Sensorik Dan Meningkatkan Fleksibilitas Nervus Ischiadicus Pada Hernia Nucleus Pulposus Lumbal. *Sport and Fitness Journal*, 6(1), 91–101. <https://doi.org/10.24843/spj.2018.v06.i01.p12>
- Rikesdas. (2013). Riset Kesehatan Dasar. *In Science*. <https://doi.org/10.1126/science.127.3309.1275>
- RS Hermina Depok, R. (2022). Statistik Harian Rawat Jalan Fisioterapi Rsu Hermina Depok.
- Sani, A., & Durahim, D. (2021). The Differences Effect Of Giving Snags And Mc Kenzie With Manual Traction And Mc Kenzie For Pain Reduction And Lumbal Disability In HNP Conditions. *Media Kesehatan Politeknik Kesehatan Makassar*, 16(1), 140. <https://doi.org/10.32382/medkes.v16i1.2052>
- Sipayung, I., Anggiat, L., & Soeparman. (2020a). Terapi Konvensional Dan Metode Mckenzie Pada Lansia Dengan Kondisi Low Back Pain Karena Hernia Nukleus Pulposus Lumbal : Studi Kasus Conventional Physiotherapy And Mckenzie Method For An Elderly Patient With Low Back Pain Caused By Lumbal Herniated Nucleus. *Jurnal Fisioterapi Dan Rehabilitasi (JFR)*, 4(2).
- Sipayung, I., Anggiat, L., & Soeparman. (2020b). Terapi konvensional dan metode mckenzie pada lansia dengan kondisi low back pain karena hernia nukleus pulposus lumbal. *ResearchGate*, August. <https://doi.org/10.33660/jfrwhs.v4i2.113>
- Tantangan, R., Bintang, S., & Ginting, S. (2021). Pengaruh Metode Mc Kenzie Exercise dan Core Stability Exercise terhadap Penurunan Nyeri pada Penderita Hernia Nucleus Pulposus. *Jurnal Riset Hesti Medan Akper Kesdam I/BB Medan*, 6(2), 107. <https://doi.org/10.34008/jurhesti.v6i2.240>
- Widyasari, O., & Wulandari, I. (2020). Penatalaksanaan Fisioterapi Pada Hernia Nucleus Pulposus (HNP) Dengan Modalitas Traksi Dan Mc. Kenzie Exercise Di Rso Prof Dr. R. Soeharso Surakarta. *Jurnal PENA* Vol.34 No.1 Edisi Maret 2020, 2507(1), 1–9. <https://doi.org/10.1016/j.solener.2019.02.027persen0Ahttps://www.golder.com/insights/block-caving-a-viable-alternative/persen0A??>
- Wijayani, N. K., Wahyudi, A., & Darmawijaya, I. P. (2022). Keseimbangan Dinamis Dengan Kecepatan Berjalan Pada Lansia Di Banjar Celuk Buruan Gianyar. Program Studi Fisioterapi, Fakultas Kesehatan, Sains, Dan Teknologi Universitas Dhyana Pura, 2(5), 2097–2104.
- Wijoyo, E. B., & Daulima, N. H. (2020). Optimalisasi Integritas Diri Melalui Terapi Kelompok Teraupetik Lansia: Studi Kasus. *Jurnal JKFT*, 5(2), 26. <https://doi.org/10.31000/jkft.v5i2.3919>
- Zahratur, A., & Priatna, H. (2019). Perbedaan Efektivitas Antara William Flexion Exercise Dan Core Stability Exercise Dalam Meningkatkan. *Jurnal Fisioterapi*, 19(1), 1–9.
- Zulvikar, J. (2016). Pengaruh Latihan Core Stability Statis (Plank dan Side Plank) dan



Core Stability Dinamis (Side Lying Hip Abduction dan Oblique Crunch) Terhadap

Megawanty, I. & Maulana, L  
*Fisiomu*. 2024, Vol 4(3) : 25-32  
DOI : <https://10.23917/fisiomu.v5i1.2370>

Keseimbangan. *Journal of Physical Education Health and Sport*, 3(2), 96–103.

